

Long-time-scale revivals in ion traps

H. Moya-Cessa*[†], A. Vidiella-Barranco[‡], J.A. Roversi[§], Dagoberto S. Freitas**^{††},
and S.M. Dutra^{‡‡}

Instituto de Física “Gleb Wataghin”, Universidade Estadual de Campinas, 13083-970 Campinas SP Brazil
(February 1, 2008)

In this contribution we investigate the interaction of a single ion in a trap with laser beams. Our approach, based on unitary transforming the Hamiltonian, allows its exact diagonalization without performing the Lamb-Dicke approximation. We obtain a transformed Jaynes-Cummings type Hamiltonian, and we demonstrate the existence of super-revivals in that system.

42.50.-p, 03.65.Bz, 42.50.Dv

There has been recently a great deal of interest in single trapped ions interacting with laser beams. This system allows the preparation of nonclassical states of the vibrational motion of the ion [1,2]. In fact, the generation of Fock, coherent, squeezed, [3] and Schrödinger-cat states [4] has been already accomplished. Besides, there are potential practical possibilities in several fields, such as precision spectroscopy [5] and quantum computation [6], for instance.

We would like to remark that the theoretical treatment of the interaction of a trapped ion with one or several laser beams constitutes a complicated problem. Normally the problem is solved in limiting situations, e.g., in the Lamb-Dicke regime, in which the ion is confined in a region much smaller than the laser wavelength. Other limiting situations concern the laser intensity; if the effective Rabi frequency Ω of the ion-laser interaction is such that $\Omega \ll \nu$, where ν is the trap frequency [7,8], we have the low-excitation regime. Otherwise, if $\Omega \gg \nu$, this corresponds to the strong-excitation regime [9].

Here we adopt a new approach to this problem. We depart from the full ion-laser Hamiltonian, and perform a unitary transformation that allows us to obtain a Jaynes-Cummings like Hamiltonian without the rotating-wave approximation (RWA). We therefore are able to analitically solve the problem, by applying the RWA, which corresponds to the regime $\Omega \approx \nu$. We should stress that our method makes possible to close the gap between the different intensity regimes, and moreover, with no restrictions on the Lamb-Dicke parameter.

We consider a single trapped ion interacting with two laser plane waves (frequencies ω_1 and ω_2), in a Raman-type configuration. Both laser beams, as usual, will be treated as classical fields propagating along the x axis, so that we have a one-dimensional problem. The corresponding scheme of levels is shown in Fig. (1). The lasers effectively drive the electric-dipole forbidden transition $|g\rangle \leftrightarrow |e\rangle$ (frequency ω_0), and we may have a detuning $\delta = \omega_0 - \omega_L$, where ($\omega_L = \omega_1 - \omega_2$). For a sufficiently large detuning Δ , the third level $|r\rangle$ may be adiabatically eliminated [10], and we end up with an effective two-level system. This situation is described by the following Hamiltonian [9]

$$\hat{H} = \hbar\nu\hat{a}^\dagger\hat{a} + \hbar\frac{\delta}{2}\sigma_z + \hbar\frac{\Omega}{2}\left(\hat{\sigma}_-e^{-i\eta(\hat{a}+\hat{a}^\dagger)} + \hat{\sigma}_+e^{i\eta(\hat{a}+\hat{a}^\dagger)}\right), \quad (1)$$

being η the Lamb-Dicke parameter, $\hat{\sigma}_+ = |e\rangle\langle g|$, ($\hat{\sigma}_- = |g\rangle\langle e|$) the usual electronic raising (lowering) operator, and $\hat{a}^\dagger$ ($\hat{a}$) the ion's vibrational creation (annihilation) operator.

By applying the unitary transformation

$$\hat{T} = \frac{1}{\sqrt{2}}\left\{\frac{1}{2}\left[\hat{D}^\dagger(\beta) + \hat{D}(\beta)\right]\hat{I} + \frac{1}{2}\left[\hat{D}^\dagger(\beta) - \hat{D}(\beta)\right]\hat{\sigma}_z + \hat{D}(\beta)\hat{\sigma}_+ - \hat{D}^\dagger(\beta)\hat{\sigma}_-\right\}, \quad (2)$$

to the Hamiltonian in Eq. (1), where $\hat{D}(\beta) = \exp(\beta\hat{a}^\dagger - \beta^*\hat{a})$ is Glauber's displacement operator, with $\beta = i\eta/2$, we obtain the following transformed Hamiltonian

*Electronic address: hmmc@inaoep.mx

‡Electronic address: vidiella@ifi.unicamp.br

§Electronic address: rovers@ifi.unicamp.br

**Electronic address: dfreitas@ifi.unicamp.br

‡‡Electronic address: dutra@rullm1.leidenuniv.nl

†Permanent address: INAOE, Coordinación de Óptica, Apdo. Postal 51 y 216, 72000 Puebla, Pue., Mexico

††Permanent address: Universidade Estadual de Feira de Santana, Departamento de Física, 44031-460, Feira de Santana, BA, Brazil

$$\hat{\mathcal{H}} \equiv \hat{T} \hat{H} \hat{T}^\dagger = \hbar\nu\hat{n} + \hbar\Omega\hat{\sigma}_z - i\hbar\frac{\eta\nu}{2} \left[(\hat{a}^\dagger - \hat{a}) - i\frac{\delta}{\eta\nu} \right] (\hat{\sigma}_- + \hat{\sigma}_+) + \hbar\nu\frac{\eta^2}{4}. \quad (3)$$

This result holds for any value of the Lamb-Dicke parameter η . Our Hamiltonian in Eq. (1) becomes a Jaynes-Cummings-type Hamiltonian, and therefore its diagonalization is allowed provided we perform the rotating wave approximation (RWA). This is possible in the regime $\nu = 2\Omega$, i.e., when the Rabi frequency of the coupling between the laser and the ion (Ω), is of the order of the vibrational frequency of the ion in the trap (ν).

Rewriting $\hat{\mathcal{H}}$ in the interaction representation,

$$\hat{\mathcal{H}} = -i\hbar\frac{\eta\nu}{2} \left[\left(\hat{a}^\dagger \hat{\sigma}_- e^{i(\nu-2\Omega)t} - h.c. \right) + \left(\hat{a}^\dagger \hat{\sigma}_+ e^{i(\nu+2\Omega)t} - h.c. \right) + -i\frac{\delta}{\eta\nu} (\hat{\sigma}_+ e^{2i\Omega t} - h.c.) \right], \quad (4)$$

we see that we may neglect the last two rapidly oscillating terms in the right hand side of Eq. (4), which corresponds to the RWA. We then obtain

$$\hat{\mathcal{H}} \approx \hat{\mathcal{H}}_{RWA} = \hbar\nu\hat{n} + \hbar\frac{\nu}{2}\hat{\sigma}_z - i\hbar g (\hat{a}^\dagger \hat{\sigma}_- - \hat{a} \hat{\sigma}_+), \quad (5)$$

which coincides with the Jaynes-Cummings Hamiltonian. The effective coupling constant is $g = \eta\nu/2$, and the term $\hbar\nu\eta^2/4$ has not been taken into account because it just represents an overall phase. We note that our final result in Eq. (5) is valid even for non-zero detunings δ .

The time evolution of the state vector, for an initial state $|\psi(0)\rangle$ is

$$|\psi(t)\rangle = \hat{T}^\dagger \hat{U}_I(t) \hat{T} |\psi(0)\rangle, \quad (6)$$

where $\hat{U}_I(t)$ is the Jaynes-Cummings evolution operator [11] in the interaction picture

$$\hat{U}_I(t) = \frac{1}{\sqrt{2}} \left\{ \frac{1}{2} [\hat{C}_{n+1} + \hat{C}_n] \hat{I} + \frac{1}{2} [\hat{C}_{n+1} - \hat{C}_n] \hat{\sigma}_z + \hat{S}_{n+1} \hat{a} \hat{\sigma}_+ - \hat{a}^\dagger \hat{S}_{n+1} \hat{\sigma}_- \right\}, \quad (7)$$

with $\hat{C}_{n+1} = \cos(gt\sqrt{\hat{a}\hat{a}^\dagger})$ and $\hat{S}_{n+1} = \sin(gt\sqrt{\hat{a}\hat{a}^\dagger})/\sqrt{\hat{a}\hat{a}^\dagger}$.

For an initial state $|\psi(0)\rangle = |e\rangle|\alpha\rangle$, or the atom in the excited state [see Fig. (1)] and the ion in a coherent (vibrational) state, the mean number of vibrational quanta as a function of the (dimensionless) scaled time $\tau = gt$, or $\langle\hat{n}\rangle = \langle\psi(\tau)|\hat{n}|\psi(\tau)\rangle$ is given by

$$\begin{aligned} \langle\hat{n}\rangle &= \frac{1}{2} \sum_{n=0}^{\infty} \left(|c_{n,e}(t)|^2 + |c_{n,g}(t)|^2 \right) \left(n + \frac{\eta^2}{4} \right) \\ &+ \left(\frac{i\eta}{4} \right) \sum_{n=0}^{\infty} \sqrt{n+1} [c_{n+1,e}^*(t) c_{n,g}(t) - c_{n,e}^*(t) c_{n+1,g}(t) - c.c.] \end{aligned} \quad (8)$$

where

$$c_{n,e}(t) = \exp - |\tilde{\alpha}|^2 / 2 \left[\frac{\tilde{\alpha}^n}{\sqrt{n!}} \cos(gt\sqrt{n+1}) + \frac{\tilde{\alpha}^{n+1}}{\sqrt{(n+1)!}} \sin(gt\sqrt{n+1}) \right], \quad (9)$$

$$c_{n,g}(t) = \exp - |\tilde{\alpha}|^2 / 2 \left[\frac{\tilde{\alpha}^{n-1}}{\sqrt{(n-1)!}} \sin(gt\sqrt{n}) - \frac{\tilde{\alpha}^n}{\sqrt{n!}} \cos(gt\sqrt{n}) \right], \quad (10)$$

and $\tilde{\alpha} = \alpha - i\eta/2$.

The structure of the equation above is similar to the one obtained for $\langle\hat{n}\rangle$ in the driven-Jaynes-Cummings model (DJCM) [12]. Therefore we expect the phenomenon of super-revivals to be present in the ion system, as an analogy to the DJCM. Super-revivals are long time scale revivals arising in the atom-field dynamics. This peculiar behavior is illustrated in Fig. (2). The super-revivals in $\langle\hat{n}\rangle$ occur for a time $\tau_{sr} \approx 4|\alpha|^2\tau_r$ ($\tau_r = 2\pi\sqrt{n}/\Omega$ is the revival time). We note that the existence or not of super-revivals is narrowly connected to the preparation of the initial vibrational state. For instance, if we have $\alpha = (5.0, 0.5)$ and $\eta = 0.5$, the super-revivals do not occur [see Fig. (2a)], and we have ordinary revivals only. However, for appropriate values of the parameters $\alpha = (0.5, 5.0)$ and $\eta = 0.5$, super-revivals take place in that system [see Fig. (2b)].

We also would like to point out that because dissipation in ion traps is much smaller than in cavities, it might be allowed the experimental observation of super-revivals in that system.

In conclusion, we have presented a novel approach for studying the dynamics of the trapped ions driven by laser beams. We have succeeded in linearizing the Hamiltonian through the application of a unitary transformation. Then we were able to treat the problem in a specific intensity regime ($\Omega \approx \nu$), without the need of performing the Lamb-Dicke approximation. As a result we have predicted the existence of long time-scale revivals (super-revivals) for the excitation number in trapped ions.




FIG. 1. Configuration of levels of the trapped ion interacting with two laser beams, of frequencies ω_1 and ω_2 .




FIG. 2. Plot of the mean excitation number $\langle \hat{a}^\dagger \hat{a} \rangle$ as a function of the (dimensionless) scaled time $\tau = gt$. a) Ordinary revivals occurring for $\alpha = (5.0, 0.5)$, and b) super-revivals occurring for $\alpha = (0.5, 5.0)$. In both cases the Lamb-Dicke parameter is $\eta = 0.5$.

One of us, H.M.-C., thanks W. Vogel for useful comments. This work was partially supported by Consejo Nacional de Ciencia y Tecnología (CONACyT), México, Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Brazil, Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Brazil, and International Centre for Theoretical Physics (ICTP), Italy.

- [1] D.J. Heinzen, and D.J. Wineland, Phys. Rev. A, **42**, 2977 (1990); J.I. Cirac, A.S. Parkins, R. Blatt, and P. Zoller, Phys. Rev. Lett., **70**, 556 (1996).
- [2] R.L. de Matos Filho and W. Vogel, Phys. Rev. Lett., **76**, 608 (1996); R.L. de Matos Filho and W. Vogel, Phys. Rev. A, **54**, 4560 (1996).
- [3] D.M. Meekhof, C. Monroe, B.E. King, W.M. Itano, and D.J. Wineland, Phys. Rev. Lett., **76**, 1796 (1996).
- [4] C. Monroe, D.M. Meekhof, B.E. King, and D.J. Wineland, *Science*, **272**, 1131 (1996).
- [5] D.J. Wineland, J.J. Bollinger, W.M. Itano, F.L. Moore, and D.J. Heinzen, Phys. Rev. A, **46**, R6797 (1992).
- [6] J.I. Cirac, P. Zoller, Phys. Rev. Lett., **74**, 4091 (1995).
- [7] C.A. Blockey, D.F. Walls, and H. Risken, Europhys. Lett. **77**, 509 (1992); J.I. Cirac, R. Blatt, A.S. Parkins, and P. Zoller, Phys. Rev. A, **49**, 1202 (1994).
- [8] J.I. Cirac, R. Blatt, A.S. Parkins, and P. Zoller, Phys. Rev. Lett., **70**, 762 (1993).
- [9] J.F. Poyatos, J.I. Cirac, R. Blatt, and P. Zoller, Phys. Rev. A, **54**, 1532 (1996).
- [10] C.C. Gerry, J.H. Eberly, Phys. Rev. A, **42**, 6805 (1990).
- [11] S. Stenholm, Phys. Rep. C, **6**, 1 (1973).
- [12] S. M. Dutra, P. L. Knight, and H. Moya-Cessa, Phys. Rev. A **49**, 1993 (1994).